

7. M Gissler et. al., “Pregnancy Associated Deaths in Finland 1987-1994 -- definition problems and benefits of record linkage,” *Acta Obstetrica et Gynecologica Scandinavica* 76:651-657, 1997; M. Gissler, “Injury deaths, suicides and homicides associated with pregnancy, Finland 1987-2000,” *European J. Public Health* 15(5):459-63, 2005 . This study looked at death rates for up to one year after the pregnancy ended. Another study found that, compared to women who gave birth, women who had abortions had a 62% higher risk of death from all causes for at least *eight* years after their pregnancies. See DC Reardon et. al., “Deaths Associated With Pregnancy Outcome: A Record Linkage Study of Low Income Women,” *Southern Medical Journal* 95(8):834-41, Aug. 2002.

8. M. Gissler et. al., “Injury deaths, suicides and homicides associated with pregnancy, Finland 1987-2000,” *European J. Public Health* 15(5):459-63, 2005; Gissler et. al., “Suicides After Pregnancy in Finland: 1987-94: register linkage study,” *British Medical Journal*, 313: 1431-1434, 1996. This looked at suicide rates for an average for up to one year after the end of the pregnancy. Another study that looked at suicide rates for up to *eight years* after the pregnancy ended found that women who aborted had a 2.5 times higher suicide rate. See DC Reardon et. al., “Deaths Associated With Pregnancy Outcome: A Record Linkage Study of Low Income Women,” *Southern Medical Journal* 95(8):834-41, Aug. 2002.

9. N. Maconochie, P. Doyle, S. Prior, R. Simmons, “Risk factors for first trimester

miscarriage—results from a UK-population-based case-control study,” *BJOG: An International Journal of Obstetrics & Gynaecology*, Dec 2006.

Part I Answers

1. d) 64% 1
2. f) homicide 2
3. b) 54% 1
4. b) 52% 1
5. d) 67% 1
6. e) 84% 1
7. e) 79% 1
8. d) 6 times as likely 3

Part II Answers

9. e) 31% 4
10. d) 10% 5
11. b) 65% higher risk 6
12. f) 65% 1
13. c) 3.5 times higher 7
14. e) 60% 1
15. a) 6 times higher 8
16. a) 60% 9

For more information — including links to published research, personal testimonies, and user-friendly resources — visit www.unchoice.info.

The Evangelization Station

Hudson, Florida, USA

E-mail: evangelization@earthlink.net

www.evangelizationstation.com

Pamphlet 639

Take the Unchoice Challenge

PART I - Before Abortion

1. What percentage of women having abortions feel pressured by others into abortion?

- (a) 12% (b) 51% (c) 27% (d) 64% (e) 15% (f) 32%

2. What is the #1 killer of pregnant women?

- (a) breast cancer (b) heart disease (c) malpractice (d) gestational diabetes (e) second-hand smoke (f) homicide

3. How many women felt uncertain at the time of their abortion?

- (a) 12% (b) 54% (c) 93% (d) 22% (e) 48% (f) 26%

4. How many women felt rushed at the time of their abortion?

- (a) 12% (b) 52% (c) 93% (d) 22% (e) 48% (f) 26%

5. How many women received no counseling by the clinic or other service-providers who performed the abortion?

- (a) 0-10% (b) 28% (c) 41% (d) 67% (e) 76% (f) 83%

6. How many women felt they did not receive adequate counseling before their abortion?

(a) 2-7% (b) 11-29% (c) 51% (d) 79% (e) 84% (f) 92%

7. How many were not informed by the abortion provider about available alternatives?

(a) 81% (b) 79% (c) 11% (d) 66% (e) 79% (f) 82%

8. If a teenager has had an abortion within the last six months, how much more likely is she to commit suicide?

(a) two times more (b) 3-4 times more (c) 5.5 times more (d) 6 times more (e) 8.2 times more (f) 15 times more

PART II - After Abortion

9. How many women having abortions report suffering physical health complications after abortion?

(a) 2.75% (b) 5% (c) 13.2% (d) 28% (e) 31% (f) 78%

10. What percentage of women having abortions suffer immediate, potentially life-threatening complications?

(a) 1% (b) 1.3% (c) 2% (d) 10% (e) 25% (f) 50%

11. What is the risk of clinical depression among women following abortion vs. childbirth?

(a) 3% higher (b) 65% higher (c) 10% higher (d) 26.2% higher (e) 5% higher (f) 73% higher

12. What percentage of women suffer symptoms of Post-Traumatic Stress Disorder (PTSD) after abortion?

(a) none (b) 6% (c) 22% (d) 42% (e) 58% (f) 65%

13. When considering death from various causes, how much higher are women's death rates after abortion vs. giving birth?

(a) about the same (b) twice as high (c) 3.5 times as high (d) 5 times higher (e) 10 times higher (f) 22 times higher

14. How many women surveyed after abortion responded that: "Part of me died."?

(a) 11.5% (b) 30% (c) 45% (d) 53% (e) 60% (f) 71%

15. How do suicide rates among women who have had abortions compare to suicide rates among those who gave birth?

(a) 6 times higher (b) twice as high (c) 4.3 times higher (d) 5 times higher (e) 8 times higher (f) 9-10 times higher

16. Abortion increases a woman's risk of future miscarriages by how much?

(a) 60% (b) 20% (c) 44% (d) 38% (e) 32% (f) 28%

References:

1. Rue et. al., "Induced abortion and traumatic stress: A preliminary comparison of American and Russian women," *Medical Science Monitor* 10(10): SR5-16, 2004.
2. I.L. Horton and D. Cheng, "Enhanced Surveillance for Pregnancy-Associated Mortality-Maryland, 1993-1998," *JAMA* 285(11): 1455-1459 (2001); J. Mcfarlane et. al., "Abuse During Pregnancy and Femicide: Urgent Implications for Women's Health," *Obstetrics & Gynecology* 100: 27-36 (2002).
3. Garfinkel et al., "Stress, Depression and Suicide: A Study of Adolescents in Minnesota," *Responding to High Risk Youth* (U. of Minnesota: Minnesota Extension Service, 1986)
4. Garfinkel et al., "Stress, Depression and Suicide: A Study of Adolescents in Minnesota," *Responding to High Risk Youth* (U. of Minnesota: Minnesota Extension Service, 1986)
5. Frank, et.al., "Induced Abortion Operations and Their Early Sequelae," *Journal of the Royal College of General Practitioners* 35(73):175-180, April 1985; Grimes and Cates, "Abortion: Methods and Complications", in *Human Reproduction*, 2nd ed., 796-813; M.A. Freedman, "Comparison of complication rates in first trimester abortions performed by physician assistants and physicians," *Am. J. Public Health* 76(5):550-554, 1986).
6. JR Cogle et. al., "Depression Associated With Abortion and Childbirth: A Long-Term Analysis of the NLSY Cohort," *Medical Science Monitor* 9(4):CR105-112, 2003.